

Addendum To: On fibre space structures of a projective irreducible symplectic manifold

Daisuke Matsushita

E-mail Address tyler@kurims.kyoto-u.ac.jp

In this note, we prove that every fibre space of a projective irreducible manifold is a lagrangian fibration.

THEOREM 1 *Let X be a projective irreducible symplectic manifold and $f : X \rightarrow B$ a fibre space with projective base B . Then f is a lagrangian fibration, that is a general fibre of f is a lagrangian submanifold.*

REMARK. Beauville proves that a Lagrangian fibration is a complete integrable system in [2, Proposition 1]. Thus, a general fibre of a fibre space of a projective irreducible symplectic manifold is an abelian variety.

REMARK. Markshevich states in [4, Remark 3.2] that there exists an irreducible symplectic manifold which has a family of non lagrangian tori. But this family does not form fibration.

PROOF OF THEOREM. Let ω be a nondegenerate two form on X and $\bar{\omega}$ a conjugate. Assume that $\dim X = 2n$. Then $\dim F = n$ [5, Theorem 2], where F is a general fibre of f . In order to prove that F is a Lagrangian submanifold, it is enough to show

$$\int_F \omega \wedge \bar{\omega} A^{n-2} = 0$$

where A is an ample divisor on X . Let H' be an ample divisor on B and $H := f^*H'$. Then

$$\int_F \omega \wedge \bar{\omega} A^{n-2} = c(\omega \bar{\omega} A^{n-2} H^n),$$

where c is a nonzero constant. We shall show $\omega \bar{\omega} A^{n-2} H^n = 0$. By [3, Theorem 4.7], there exists a bilinear form q_X on $H^2(X, \mathbb{C})$ which has the following properties:

$$a_0 q_X(D, D)^n = D^{2n} \quad D \in H^2(X, \mathbb{C}).$$

We consider the following equation,

$$a_0 q_X(\omega + \bar{\omega} + sA + tH, \omega + \bar{\omega} + sA + tH)^n = (\omega + \bar{\omega} + sA + tH)^{2n}.$$

Calculating the left hand side, we obtain

$$a_0(q_X(\omega + \bar{\omega}) + s^2 q_X(A) + 2s q_X(\omega + \bar{\omega}, A) + 2t q_X(\omega + \bar{\omega}, H) + 2st q_X(A, H))^n.$$

Since $\omega \in H^0(X, \Omega_X^2)$ and $A, H \in H^1(X, \Omega_X^1)$,

$$q_X(\omega + \bar{\omega}, A) = q_X(\omega + \bar{\omega}, H) = 0$$

by [1, Théorème 5]. Thus we can conclude that $\omega \bar{\omega} A^{n-2} H^n = 0$ by comparing $s^{n-2} t^n$ term of both hands sides. Q.E.D.

References

- [1] A. Beauville, *Variétés kähleriennes dont la première classes de Chern est nulle*, J. Diff. Geom., **18** (1983), 755–782.
- [2] ———, *Systèmes hamiltoniens complètement intégrables associés aux surfaces K3*, in *Problems in the theory of surfaces and their classification* Sympos. Math., XXXII, Academic Press, London, (1991), 25–31.
- [3] A. Fujiki, *On the de Rham Cohomology Group of a Compact Kähler Symplectic Manifold*, in *Algebraic geometry, Adv. Stud. Pure Math.*, vol **10**, (ed. T. Oda), Kinokuniya and North-Holland (1987), 105–165.
- [4] D.G. Markushevich, *Integrable symplectic structures on complex manifolds*, Math. USSR-Sb., **59**:2 (1988), 459–469.
- [5] D. Matsushita, *On fibre space structures of a projective irreducible symplectic manifold*, Topology, **38** (1999), 79–81.

Research Institute of Mathematical Science,
Kyoto University.

KITASHIRAKAWA, OIWAKE-CHO, KYOTO, 606-01, JAPAN.